ROBERT C. GRUMET, MD

TOTAL SHOULDER AND HEMIARTHROPLASTY PROTOCOL

Name			Date
Diagnosi			
Date of S	Surgery		
Frequenc	cy: 1 2 3 4 times/week Du	uration: 1 2 3 4 5 6 We	eeks
•		of 6 weeks. The subscapularis tender it to grown back into the humerulat side; ABD max 75° without rotate at side; ABD max 75° without rotate extension until 12 weeks post-op	
•	Weeks 6-12: Begin AAROM → AROM for internal Goals: Increase ROM as tolerated with Begin light resisted ER/FF/ABD: isom No resisted internal rotation/backwards No scapular retractions with bands yet	h gentle passive stretching at end ra tetrics and bands, concentric motion s extension until 12 weeks post-op	inges
•	Months 3-12: Begin resisted IR/BE (isometrics/bands Advance strengthening as tolerated; 10 Increase ROM to full with passive stre Begin eccentric motions, plyometrics,	reps/1 set per exercise for rotator of tching at end ranges	cuff, deltoid, and scapular stabilizers.
Commer	nts:		
Fun	ctional Capacity EvaluationWo	rk Hardening/Work Conditioning	Teach HEP
Modaliti Elect	es tric StimulationUltrasound I	ontophoresisPhonophoresis	TENS Heat before/after
Ice b	efore/afterTrigger points massage	Other	Therapist's discretion
Signatur	e	Date	

Orthopaedic Specialty Institute 280 S. Main Street, Suite 200 Orange, CA. 92868 Main: (714) 634-4567 Fax: (714) 634-4569